

			Disorder in parents										
			None	Psychosis	Bipolar disorder	Depression	Anxiety disorders	Substance use disorders	Borderline pers. disorder	ADHD	Disruptive disorders	OCD	Eating disorders
Disorder in offspring	Psychosis	(followed until adulthood)	2% (1-4%) (k=13,n=1801)	21% (12-36%) (k=10,n=901)	4% (3-7%) (k=3,n=471)	4% (1-12%) (k=2,n=197)		5% (2-14%) (k=1,n=74)					33% (13-62%) (k=1,n=12)
	Bipolar disorder	(followed until adulthood)	2% (1-4%) (k=12,n=2199)	2% (0-11%) (k=2,n=79)	13% (10-17%) (k=11,n=1721)	7% (4-13%) (k=5,n=1580)							
	Depression	(followed until adolescence or adulthood)	11% (9-14%) (k=36,n=7937)	15% (9-24%) (k=6,n=532)	26% (20-33%) (k=14,n=1958)	34% (26-43%) (k=18,n=4891)	12% (7-21%) (k=2,n=84)	15% (10-23%) (k=4,n=675)			21% (11-36%) (k=1,n=43)		43% (33-53%) (k=1,n=96)
	Anxiety disorders	(followed until adolescence or adulthood)	16% (13-19%) (k=28,n=5899)	12% (6-21%) (k=4,n=137)	27% (20-36%) (k=13,n=1880)	29% (22-37%) (k=13,n=2976)	31% (7-72%) (k=2,n=266)	17% (9-32%) (k=4,n=539)			51% (37-66%) (k=1,n=43)		29% (21-39%) (k=1,n=96)
	Substance use disorders	(followed until adulthood)	14% (10-18%) (k=22,n=5489)	16% (7-34%) (k=4,n=297)	21% (15-29%) (k=8,n=1151)	34% (22-49%) (k=8,n=1842)		35% (10-73%) (k=5,n=762)					48% (38-58%) (k=1,n=96)
	Borderline pers. disorder	(followed until adulthood)	17% (6-37%) (k=1,n=24)	9% (1-44%) (k=2,n=176)									
	ADHD	(any follow-up)	8% (6-11%) (k=23,n=2964)	32% (12-60%) (k=2,n=240)	14% (9-20%) (k=14,n=2077)	11% (7-16%) (k=9,n=1009)	8% (3-21%) (k=2,n=266)	16% (13-20%) (k=5,n=423)			23% (13-38%) (k=1,n=43)		
	Disruptive disorders	(any follow-up)	4% (3-7%) (k=16,n=2615)	7% (4-11%) (k=3,n=246)	13% (7-23%) (k=8,n=1294)	11% (3-30%) (k=4,n=231)	8% (3-21%) (k=2,n=266)	14% (11-18%) (k=4,n=360)			15% (7-29%) (k=1,n=41)		11% (6-20%) (k=1,n=96)
	OCD	(any follow-up)	1% (1-2%) (k=16,n=2514)	1% (0-16%) (k=1,n=41)	3% (1-5%) (k=8,n=1382)	3% (2-6%) (k=5,n=1586)	11% (4-29%) (k=1,n=27)	2% (0-6%) (k=1,n=124)			23% (13-38%) (k=1,n=43)		
	Eating disorders	(followed until adolescence or adulthood)	3% (1-7%) (k=4,n=368)	3% (1-12%) (k=2,n=60)	1% (0-2%) (k=3,n=428)						7% (2-20%) (k=1,n=43)		
	Any mental disorder	(followed until adulthood)	30% (23-38%) (k=13,n=2169)	56% (41-71%) (k=5,n=596)	66% (53-78%) (k=4,n=275)	75% (52-90%) (k=3,n=287)		55% (44-66%) (k=1,n=74)					83% (75-89%) (k=2,n=108)